

No association between telomere length-related loci and number of cutaneous nevi

Xin Li¹, Geyu Liang², Mengmeng Du³, Immaculata De Vivo^{1,4}, Hongmei Nan⁵

¹Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA

²Key Laboratory of Environmental Medicine Engineering, Ministry of Education, School of Public Health, Southeast University, Nanjing, Jiangsu, China

³Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY, USA

⁴Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA

⁵Department of Epidemiology, Richard M. Fairbanks School of Public Health, Indiana University, and Indiana University Melvin and Bren Simon Cancer Center, Indianapolis, IN, USA

Correspondence to: Hongmei Nan, **email:** hnan@iu.edu

Keywords: telomere length-related loci, melanoma, nevus count, genome-wide association study, genetic score

Received: July 14, 2016

Accepted: September 07, 2016

Published: September 14, 2016

ABSTRACT

Background: Longer telomeres have been associated both with increased melanoma risk and increased nevus counts. Nevus count is one of the strongest risk factors for melanoma. Recent data showed that a genetic score derived by telomere length-related single nucleotide polymorphisms (SNPs) was strongly associated with melanoma risk; however, the relationships between these SNPs and number of cutaneous nevi have not been investigated.

Methods: We evaluated the associations between telomere length-related SNPs reported by previous genome-wide association study (GWAS) and nevus counts among 15,955 participants of European Ancestry in the Nurses' Health Study and Health Professionals Follow-up Study.

Results: None of the SNPs was associated with nevus counts, nor was the genetic score combining the dosage of alleles related to increased telomere length.

Conclusions: The telomere length-related SNPs identified by published GWAS do not appear to play an important role in nevus formation. Genetic determinants of telomere length reported by GWAS do not explain the observed epidemiologic association between telomere length and nevus counts.

INTRODUCTION

Telomeres are long hexameric (TTAGGG)_n repeats at the ends of linear eukaryotic chromosomes that shorten after each round of DNA replication. They play a critical role in maintaining genomic stability [1]. Nevi occur when melanocytes cluster, and the number of melanocytic nevi in sun-exposed sites is a strong risk factor for cutaneous melanoma [2]. Han and colleagues found that shorter telomeres were associated with both a lower risk of melanoma and a decreased nevus count [3, 4]. In addition, positive associations between telomere length (TL) and nevi number and size have been reported previously [5]. It has been hypothesized that these phenotypes share common genetic components. In a recent study by Iles *et al.* [6]. The authors discovered a highly statistically

significant association between a genetic score derived by TL-related SNPs and risk of melanoma (P -value = 8.92×10^{-9}). However, it is unclear whether these genetic variants also influence the number of cutaneous nevi. We therefore evaluated the relationship between TL-related single nucleotide polymorphisms (SNPs) and nevus count.

RESULTS

We did not observe significant associations between any of the seven TL-related SNPs and nevus count (Table 1). Mean values and ranges of the genetic scores combining all 7 independent SNPs were similar among the Illumina, Affy, and Omni datasets. We found that genetic scores were not significantly associated with nevus count in our data. The results for simple count score

Table 1: Results for each telomere length-related SNP, including their associations with melanoma risk and nevus count

SNP	Chr	Position	Related gene	Minor allele	MAF in telomere GWAS [13]	MAF in our data	Average imputation R-square ¹	Melanoma beta ²	Melanoma P value ²	Nevus count beta	Nevus count P value
rs10936599	3	169492101	<i>TERC</i>	T	0.252	0.241	0.993	-0.079	0.0003	-0.020	0.48
rs2736100	5	1286516	<i>TERT</i>	C	0.486	0.506	0.988	0.078	0.02	-0.043	0.08
rs7675998	4	164007820	<i>NAF1</i>	A	0.217	0.222	0.988	-0.063	0.03	-0.036	0.22
rs9420907	10	105676465	<i>OBFC1</i>	C	0.135	0.142	1.000	0.083	0.001	0.051	0.14
rs8105767	19	22215441	<i>ZNF208</i>	G	0.291	0.290	0.996	0.028	0.16	0.025	0.34
rs755017	20	62421622	<i>RTEL1</i>	G	0.131	0.123	0.952	0.026	0.35	0.004	0.92
rs11125529	2	54475866	<i>ACYP2</i>	A	0.142	0.134	0.999	-0.004	0.86	0.020	0.56

1 Average imputation R square quality metric of Illumina, Affy, and Omni datasets;

2 Information on associations between SNPs and risk of melanoma was obtained from paper by Iles et al. [6].

Table 2: Associations between genetic scores (GSs) of telomere length-related SNPs and nevus count¹

	Illumina			Affy			Omni			Meta-analysis ²		
	Beta	SE	P-value	Beta	SE	P-value	Beta	SE	P-value	Beta	P-value	P Heterogeneity
Simple count GS	0.003	0.008	0.67	0.009	0.007	0.21	-0.001	0.008	0.90	0.004	0.33	0.66
Weighted GS	0.004	0.008	0.64	0.009	0.007	0.21	-0.004	0.009	0.64	0.005	0.42	0.50

1 Linear regression models were used to assess the relationship between genetic scores (rescaled) and nevus count, adjusting for age, gender, and the top three principle components;

2 Analyses were first conducted within each of the platform-specific genetic datasets. We used fixed-effect meta-analysis to obtain a combined estimate.

(beta = 0.004, *p*-value = 0.33) and weighted score (beta = 0.005, *p*-value = 0.42) were similar (Table 2). We calculated the power using the pwr package for multiple linear regression in R; our data had 99% power at two-sided alpha of 0.05 to detect an effect size beta of 0.004.

DISCUSSION

Despite epidemiologic evidence of a link between TL and nevus count, and the strong association between TL-related SNPs and melanoma, we did not detect an association between TL GWAS SNPs and number of nevi. This is consistent with findings of some experimental studies that telomere attrition may not be involved in the initial formation of melanocytic nevi, because telomere length in nevi did not differ significantly from surrounding tissue and nevi generally lack expression of p53 and p21, markers of telomere-induced senescence [7]. In addition, both these phenotypes have long been recognized as the result of combined genetic and environmental risk factors. Thus, the reported association might be largely due to shared environmental risk factors (such as UV exposure) or regulations beyond the gene level. Furthermore, it is possible that the selected SNPs do not cover those genes involved in both telomere

length and nevus count, as our knowledge of TL-related loci remains incomplete. In summary, we did not find evidence of associations between the selected TL-related alleles and nevus count. This suggests that the effect of telomere genetic score on melanoma risk may not be mediated through nevus counts. Future work is needed to screen more novel TL-related genetic variants, and investigate the association between these phenotypes at the level of other “-omics”.

MATERIALS AND METHODS

Eighteen case-control studies nested within the Nurses' Health Study (NHS) and the Health Professionals Follow-up Study (HPFS) were combined into three compiled datasets based on their genotype platforms: Affymetrix (Affy), Illumina HumanHap series (Illumina), or Illumina Omni Express (Omni) (Supplementary Tables S1 and S2). Detailed descriptions on study population, genotyping procedure, quality control and imputation are provided in Supplementary Materials.

In total, 15,955 participants of European ancestry were included in the current study. In both the NHS and HPFS, participants were asked to provide information on melanocytic nevus count (larger than 3 mm in diameter)

on arms by choosing from the following categories: 1 = none, 2 = 1–2, 3 = 3–5, 4 = 6–9, 5 = 10–14, 6 = 15–20, and 7 = 21+. Previous studies have evaluated the validity of self-reported information on nevus counts. The majority of studies on nevus counts have shown substantial agreement between nevus self-counts and dermatologist counts [8, 9]. The Spearman's correlation coefficient was 0.91 between the two measurements [10]. In our cohorts, the ordinal variable of nevus count is a highly significant predictor of melanoma risk [11]. A genome-wide association study on nevi number using our genetic datasets has revealed biologically plausible susceptibility loci for nevus count and melanoma risk, and the same loci were replicated by other independent studies. In the current study, we used number of nevi on the arms as proxy for the total body nevus counts, which has been proven to be acceptable [12].

We obtained seven independent TL-related SNPs from the largest genome-wide association study (GWAS) on TL [13]. The same SNPs were examined by Iles *et al.* [6] for the association with melanoma risk. For each participant, we summed the dosage of alleles related to increase in TL of the seven SNPs to obtain the simple count genetic score. We also constructed a weighted score by multiplying the dosage of effect alleles by the corresponding regression coefficients in the original GWAS paper [13] and then summing the products. Both the simple count score and the weighted score were rescaled to a mean of 14 alleles (2 alleles * 7 SNPs) before testing their associations with nevus count to make results comparable. We presented formula and summary statistics of the genetic scores in Supplementary Table S3.

We regressed nevus counts (0 = none, 1.5 = 1–2, 4 = 3–5, 7.5 = 6–9, 10 = 10+) on the dosage of each SNP and the genetic scores, respectively, adjusted for age, sex, and the top three principle components (PCs). All the analyses were first conducted within each of the platform-specific datasets, and then combined by inverse-variance-weighted meta-analysis if results were not significantly different. ProbABEL package and R-3.0.2 were used to perform these tests. We considered 2-sided *P* values less than 0.05 to be statistically significant.

ACKNOWLEDGMENTS

We are indebted to the participants in the NHS and HPFS for their dedication to this research. We thank the following state cancer registries for their help: Alabama, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Virginia, Washington, and Wyoming. The authors assume full responsibility for analyses and interpretation of these data.

CONFLICTS OF INTEREST

The authors state no conflicts of interest.

GRANT SUPPORT

This work is supported by NIH R01 CA49449, P01 CA87969, UM1 CA186107, UM1 CA167552, and P30 CA008748.

REFERENCES

1. Stewart SA, Weinberg RA. Telomeres: cancer to human aging. *Annu Rev Cell Dev Biol.* 2006; 22:531–557.
2. Gandini S, Sera F, Cattaruzza MS, Pasquini P, Abeni D, Boyle P, Melchi CF. Meta-analysis of risk factors for cutaneous melanoma: I. Common and atypical naevi. *European Journal of Cancer.* 2005; 4:28–44.
3. Han J, Qureshi AA, Prescott J, Guo Q, Ye L, Hunter DJ, De Vivo I. A prospective study of telomere length and the risk of skin cancer. *Journal of Investigative Dermatology.* 2009; 12:415–421.
4. Nan H, Du M, De Vivo I, Manson JE, Liu S, McTiernan A, Curb JD, Lessin LS, Bonner MR, Guo Q. Shorter telomeres associate with a reduced risk of melanoma development. *Cancer research.* 2011; 7:6758–6763.
5. Bataille V, Kato BS, Falchi M, Gardner J, Kimura M, Lens M, Perks U, Valdes AM, Bennett DC, Aviv A. Nevus size and number are associated with telomere length and represent potential markers of a decreased senescence *in vivo*. *Cancer Epidemiology Biomarkers & Prevention.* 2007; 1:1499–1502.
6. Iles MM, Bishop DT, Taylor JC, Hayward NK, Brossard M, Cust AE, Dunning AM, Lee JE, Moses EK, Akslen LA. The effect on melanoma risk of genes previously associated with telomere length. *Journal of the National Cancer Institute.* 2014; 10:dju267.
7. Mooi W, Peeper D. Oncogene-induced cell senescence—halting on the road to cancer. *New England Journal of Medicine.* 2006; 35:1037–1046.
8. Buettner PG, Garbe C. Agreement between self-assessment of melanocytic nevi by patients and dermatologic examination. *American journal of epidemiology.* 2000; 15:72–77.
9. Little P, Keefe M, White J. Self screening for risk of melanoma: validity of self mole counting by patients in a single general practice. *Bmj.* 1995; 31:912–916.
10. Mikkilineni R, Weinstock MA. Is the self-counting of moles a valid method of assessing melanoma risk? *Archives of dermatology.* 2000; 13:1550–1551.
11. Cho E, Rosner BA, Feskanich D, Colditz GA. Risk factors and individual probabilities of melanoma for whites. *Journal of clinical oncology.* 2005; 2:2669–2675.

12. Bain C, Colditz GA, Willett WC, Stampfer MJ, Green A, Bronstein BR, Mihm MC, Rosner B, Hennekens CH, Speizer FE. Self-reports of mole counts and cutaneous malignant melanoma in women: methodological issues and risk of disease. *American journal of epidemiology*. 1988; 12:703–712.
13. Codd V, Nelson CP, Albrecht E, Mangino M, Deelen J, Buxton JL, Hottenga JJ, Fischer K, Esko T, Surakka I. Identification of seven loci affecting mean telomere length and their association with disease. *Nature genetics*. 2013; 4:422–427.