

Identification of *RECK* as an evolutionarily conserved tumor suppressor gene for zebrafish malignant peripheral nerve sheath tumors

Rashmi Kumari^{1,*}, Martin R. Silic^{1,*}, Yava L. Jones-Hall^{1,2}, Alexandra Nin-Velez¹, Jer-Yen Yang^{3,4,5}, Suresh K. Mittal^{1,2,4} and GuangJun Zhang^{1,2,4,5}

¹Department of Comparative Pathobiology, Purdue University, West Lafayette, Indiana 47907, USA

²Purdue Institute for Inflammation, Immunology and Infectious Disease (PI4D), Purdue University, West Lafayette, Indiana 47907, USA

³Department of Basic Medical Sciences, Purdue University, West Lafayette, Indiana 47907, USA

⁴Purdue University Center for Cancer Research, Purdue University, West Lafayette, Indiana 47907, USA

⁵Purdue Institute for Integrative Neuroscience, Purdue University, West Lafayette, Indiana 47907, USA

*These authors contributed equally to this work

Correspondence to: GuangJun Zhang, **email:** gjzhang@purdue.edu

Keywords: *RECK*; tumor suppressor gene (*TSG*); *MPNST*; *pPNET*; zebrafish

Abbreviations: *MPNST*: malignant peripheral nerve sheath tumor; *pPNET*: peripheral primitive neuro-ectodermal tumor; *TSG*: tumor suppressor gene; *CNAs*: copy number alterations; *LOH*: loss of heterozygosity

Received: May 16, 2017

Accepted: April 08, 2018

Published: May 04, 2018

Copyright: Kumari et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Malignant peripheral nerve sheath tumors (MPNSTs) are a type of sarcoma with poor prognosis due to their complex genetic changes, invasive growth, and insensitivity to chemo- and radiotherapies. One of the most frequently lost chromosome arms in human MPNSTs is chromosome 9p. However, the cancer driver genes located on it remain largely unknown, except the tumor suppressor gene, p16 (*INK4*)/*CDKN2A*. Previously, we identified *RECK* as a tumor suppressor gene candidate on chromosome 9p using zebrafish-human comparative oncogenomics. In this study, we investigated the tumorigenesis of the *reck* gene using zebrafish genetic models in both *tp53* and *ribosomal protein* gene mutation background. We also examined the biological effects of *RECK* gene restoration in human MPNST cell lines. These results provide the first genetic evidence that *reck* is a *bona fide* tumor suppressor gene for MPNSTs in zebrafish. In addition, restoration of the *RECK* gene in human MPNST cells leads to growth inhibition suggesting that the reactivation of *RECK* could serve as a potential therapeutic strategy for MPNSTs.

INTRODUCTION

Malignant peripheral nerve sheath tumors (MPNSTs) are a type of sarcoma that originate from the neural crest/Schwann cell lineage in humans, and have a poor prognosis due to their complex genetic changes, invasive growth nature, and insensitiveness to chemo- and radiotherapies [1–3]. For the etiology of MPNST, only a few cancer genes are currently known to be related to this type of malignancy: *NF1*, *NF2*, *SMARCB1* and *LZTR1*

[4, 5]. Surgical resection is the mainstay of treatment, but often compromised by many factors such as distant metastasis, local damage, and inoperable locations [6]. Moreover, there is currently no targeted therapy available for this type of malignancy, mainly due to the lack of knowledge of its cancer driver genes.

Cancer driver genes are a small number of genes found mutated in cancer genomes, and their mutations biologically contribute to cancer initiation and progression [7]. The remaining mutated genes in the cancer genome

are defined as passenger genes, if their mutations have no impact on tumorigenesis. Based on gain or loss of function, cancer driver genes can be divided into either oncogenes or tumor suppressor genes (TSGs). One of the goals of current cancer research is to identify all the cancer driver genes, so that they can be pursued as new targets for the development of cancer therapy. Targeted cancer therapy has emerged as a better alternative to other treatment options because of its high specificity and fewer side effects. Effective targeted cancer therapies require accurate identification of the targetable cancer driver genes.

Identification of cancer driver genes is difficult, due to the plethora of mutations in the cancer cell genomes and the time required for functional validation. One of the extreme situations is represented with aneuploid chromosomes, which are found in more than 90% of solid tumors and 75% of hematopoietic tumors [8]. This difficulty stems from the large number of genes on the chromosome arm. Even if the driver genes have no mutations, it is not possible to distinguish driver from passenger genes despite advances in modern technologies, such as parallel sequencing and microarray. Cross-species comparative oncogenomics emerged as a solution to pinpoint cancer driver candidates, as the cancer driver genes tend to be enriched in common genetic mutations, due to the gene functional conservation. This approach has already been demonstrated to be effective in mice and dogs [9–12]. The research community has recently showed that zebrafish-human comparative oncogenomics is particularly effective with large aneuploid chromosomes because gene synteny was extensively re-organized during vertebrate evolution [13–15]. Furthermore, the zebrafish model provides a valuable system for functional studies of candidate drivers from omics studies. It has been well-established that zebrafish tumors share many similarities with humans including histology, etiology of genetic mutations (*tp53*, *pten*, etc.), the transcriptome, and copy number alterations (CNAs) [16, 17].

Chromosome 9p is one of the most frequently lost chromosome arms in human MPNSTs and many other types of solid tumors [18–24]. *CDKN2A/p16/INK4* is one of the currently known TSGs on this chromosome arm, but this does not exclude the existence of other TSGs on 9p. Using zebrafish-human comparative oncogenomics, on human chromosome 9p, we have identified the *RECK* gene as a cancer driver candidate that could potentially be a tumor suppressor gene for MPNSTs [13]. *RECK* is located on chromosome 9p, which is frequently lost in human MPNSTs [25]. The *RECK* (reversion-inducing cysteine-rich protein with Kazal motifs) gene encodes a 110 kDa membrane-bound glycoprotein [26]. It was discovered as a gene that was able to rescue the aberrant cell morphology of RAS-transformed fibroblasts [27]. The *Reck* gene is expressed in many mouse tissues, and it was found to be down-regulated in many types of human cancers such as the liver, pancreatic, breast, colon, lung, melanoma, and fibrosarcoma [26, 28]. Low expression of

RECK was reported to correspond to biological malignancy, such as invasiveness and metastasis [26, 29, 30]. Thus, the *RECK* gene was believed to be a tumor suppressor gene, but *in vivo* experimental evidence was missing to support this hypothesis. Knockout mice indicate that *Reck* is a developmentally essential gene and is critical for angiogenesis through regulating matrix metalloproteinases [31]. These mouse embryos die at stage E10.5 in the homozygous knockouts, due to hemorrhage and neuronal defects [31]. Moreover, there is no report of spontaneous tumors in the heterozygous *RECK* knockout mice. Whether the *Reck* gene functions as a TSG *in vivo* remains unknown, although it is worth noting that evidence from cell culture models uphold this hypothesis [26].

Here, we performed functional genetic studies on the zebrafish *reck* gene, and provided the first evidence that zebrafish *reck* is a tumor suppressor gene *in vivo*.

RESULTS

Zebrafish are becoming a popular cancer model, due to their tractable genetics and conserved vertebrate biology similar to humans [16]. In zebrafish, MPNSTs can be initiated with *tp53* homozygous point mutations [32], or heterozygous ribosomal protein gene (*rp*) mutations [33]. Previously, we have found that zebrafish and human MPNSTs share high similarities, such as histology, aneuploidy, and copy number alterations [13, 14, 32, 33]. To investigate whether *RECK* is a *bona fide* tumor suppressor gene in MPNSTs, we took advantage of the zebrafish genetic model. First, we re-analyzed our previous DNA copy number data from both zebrafish and human MPNSTs [13]. In both human and zebrafish MPNSTs, about 39% of tumors lose DNA copy numbers of the *RECK* gene (Figure 1A–1B). As the zebrafish genome usually contains duplicates for some of the genes, due to teleost whole genome duplication, we performed a BLAT search using the human *RECK* gene as a bait in both NCBI and Ensembl. Both searches suggested that a single copy of the *reck* gene is present in the zebrafish genome. The zebrafish *Reck* shares 65% protein sequence similarity with human *RECK*, both proteins are coded by 21 exons (Figure 1C), and both feature a KAZAL_FS (Kazal type serine protease inhibitor and follistatin-like) domain. To further demonstrate the zebrafish *reck* gene identity, we conducted phylogenetic analysis using protein sequences from representative species. Our maximum likelihood phylogenetic analysis revealed that *RECK* is a conserved gene in metazoans (Figure 1D), indicating that zebrafish *reck* may have similar biological functions with the human *RECK* gene.

The zebrafish *reck* gene is found to be expressed in the developing neural crest cells, and is required for appropriate sensory neuron formation in a cell-autonomous fashion [34]. Since MPNST is a type of malignancy of Schwann cells that originate from the neural crest, we first validated the expression of the zebrafish *reck* gene by *In Situ* hybridization




Figure 1: Frequent loss of DNA copy numbers of the *RECK* gene in human and zebrafish MPNSTs. (A) Heat map of human chromosome 9 is shown. *RECK* is located on chromosome 9p, which is found to be under-represented in 39% (24 out of 61) of human MPNSTs [13]. The green line represents the *RECK* gene locus, and the vertical black line indicates the centromere of chromosome 9. The human *RECK* gene location and reference genome assembly are labeled above the heat map. (B) Zebrafish *reck* is located on chromosome 24, which is under-represented in about 39% (58 out of 147) of zebrafish MPNSTs. Samples are sorted top-to-bottom by decreasing deletion amplitude at their respective *RECK/reck* locus, indicated by a green line. Blue and red bars at the right side of each panel indicate samples with *RECK/reck*, losses (blue) or gains (red). Color densities are corresponding to the degree of loss and gain as previously described [13]. The zebrafish *reck* gene location and reference genome assembly are labeled above the heat map. (C) Human and zebrafish *RECK* have different intron-exon structures, but both are composed of 21 exons. The solid red/orange vertical boxes represent the gene coding regions, and the empty boxes indicate un-transcribed regions. Red/orange thin lines indicate the introns. Human *RECK* transcript: *RECK*-001 ENST00000377966.3. Zebrafish *reck* transcript: *reck*-001 ENSDART00000129135.2. (D) Maximum likelihood phylogeny of *RECK* proteins, as obtained with JTT plus gamma distances ($\alpha = 0.941$). Numbers around each node indicate bootstrap values based on 1,000 replicates. Branch lengths are proportional to expected replacements per site. The tree was rooted with placozoa (*Trichoplax adhaerens*).

during zebrafish embryogenesis. Indeed, as reported, we confirmed that *reck* is expressed in the developing neural crest cells (Supplementary Figure 1), suggesting it has a developmental role in the neural crest-derived cells.

A TSG is defined as a gene whose inactivation leads to tumor formation, and promotes tumor development *in vivo*. To functionally evaluate whether *reck* is a *bona fide* TSG in zebrafish MPNSTs, we took advantage of the currently available *reck* loss-of-function mutant, *sensory deprived* (*sdp*) [34]. If a TSG is strong enough, we can expect the development of spontaneous tumors in the loss-of-function mutant animals in a relatively short time. However, if a TSG is weak, we may need a longer follow-up time for tumorigenesis requiring a large number of animals. Based on the principle of genetic synthetic/additive effect, we hypothesized that loss of *reck* will accelerate tumorigenesis in a *tp53* or *ribosomal protein* mutant background, as both types of mutation were known to cause spontaneous MPNSTs in zebrafish [32, 33].

To test this hypothesis, we crossed the *reck* mutant, *sdp*^{w12/+}, with the *tp53* or *ribosomal protein L35* (*rpL35*) mutant, and tracked the development of tumors. In zebrafish, 17 *rp* gene mutants were known to cause MPNSTs and the impact of these genetic mutations was believed to be similar [33, 35, 36]. Here, we chose the *rpL35* mutant, hi258, as a representative of the *rp* TSGs. As expected, the *reck* mutation accelerated tumor formation rate in both mutant backgrounds (Figure 2A–2B), demonstrating that *reck* is indeed a *bona fide* TSG. Interestingly, we noticed that there was a tumor spectrum shift in the *reck* and *rpL35* fish cross (Figure 2C). Although most tumors were MPNSTs (Figure 2D–2E), 25% (6 out of 24) of tumors were peripheral primitive neuroectodermal tumors (pPNETs, Figure 2F–2G) in *sdp*^{w12/+};*rpL35*^{hi258/+}, while only 7% (2 out of 28) were pPNETs in *rpL35*^{hi258/+}, suggesting that the *reck* gene may have a cell type differential role during tumor formation. As we did not detect any visible tumors in the *reck* heterozygotes (*sdp*^{w12/+}) in either cross, we reasoned that *reck* may function as a TSG in a two-hit paradigm manner, exemplified by the *RBI* gene [37]. To examine this possibility, we investigated the loss of heterozygosity (LOH) in a few tumor-tail paired tissue samples. Indeed, we found 4 out of 8 cases where the wildtype allele was lost in these tumors (Figure 2H). Altogether, our results supported the hypothesis that *reck* is a TSG that may follow the two-hit paradigm of tumor suppression.

Since the DNA copy number of *RECK* is frequently lost in human MPNSTs, we hypothesized that protein levels of RECK in human MPNST cell lines are generally decreased. To test this, we performed Western blots and found that RECK was not detectable or expressed at a lower level in human MPNST cells, compared to the benign neurofibroma cell line, HEI193 (Figure 3A). TSG restoration was found to inhibit tumor growth and reverse other cancer characteristics [38], therefore, we reasoned

that restoration of *RECK* in human MPNST cells would inhibit cell growth. To demonstrate this, we first generated three tetracycline-inducible stable human MPNST cell lines using an untagged full-length *RECK* (pSLIK-neo-*RECK*) construct (Figure 3B). We chose STS26T (*TP53* mutant), S462 (*NF1* mutant), and 90-8TL (*NF1* microdeletion) human MPNST cell lines, as *TP53* and *NF1* are commonly mutated in human MPNSTs [39, 40]. Moreover, we previously found that doxycycline has no impact on the growth of these cell lines [41]. Next, we examined cell growth by MTT, colony formation assays, and soft agar anchorage-independent growth assays. Indeed, upon restoration of RECK expression the cell growth rates were decreased in all three MPNST cell lines (Figure 3C, 3F, 3I). Similarly, growth inhibition effects were also observed in the colony formation assay (Figure 3D, 3G, 3J) and the soft agar anchorage-independent growth assay (Figure 3E, 3H, 3K), suggesting that *RECK* functions as a TSG in human MPNST cells. Notably, this growth inhibition is not dependent on TP53 or NF1, suggesting that RECK is a mechanistically independent TSG.

DISCUSSION

The *RECK* gene is one of the TSG candidates on human chromosome 9p in MPNSTs and other types of tumors [18–24]. Although, accumulating evidence from cellular experiments indicates that the *RECK* gene is a TSG *in vivo*, there is no clear genetic evidence to prove this notion, since this gene is a developmentally essential gene in mouse [26, 31]. Here, our results provide the first *in vivo* evidence that *reck* is a *bona fide* tumor suppressor gene in zebrafish MPNSTs. Moreover, we validate that the restoration of RECK leads to inhibition of human MPNST cell growth.

In our genetic experiments, the loss of one *reck* gene allele indeed accelerated tumorigenesis initiated with *tp53*^{-/-}, or *rpL35*^{hi258/+}, although we did not find spontaneous tumors in zebrafish *reck* mutant, *sdp*^{w12/+}, within 2 years. These genetic synthetic/additive effects between *reck* and *tp53* or *rpL35* suggested that *reck* is a TSG and it is not in the same genetic pathway of *tp53* or *rpL35*, *i.e.* there is no direct genetic interactions. It is worth to note that the heterozygous *reck*^{sdpw12/+} alone did not lead to any visible tumors in our studies, indicating that LOH is required for *reck* tumorigenicity, or that *reck* is a weaker haploinsufficient TSG, compared to *tp53* or *rpL35* in zebrafish. TSGs usually function through LOH (two wildtype alleles are inactivated) or haploinsufficiency (one wildtype allele is inactivated) [37]. Half of our examined tumor samples (4 out of 8) showed loss of the wildtype allele, suggesting that the *reck* gene may fit LOH, and it likely follows the two-hit paradigm of tumor suppression in zebrafish. However, we cannot exclude the possibility of the haploinsufficiency mechanism, since we were only able to examine 8 samples. Further studies with more

tumors samples will be needed to address this question, and whether human RECK follows the same rule.

One of the interesting observations is that the tumor spectrum shifts in the *reck* mutant zebrafish. Generally, *tp53* or *rpL35* mutations lead to MPNSTs in zebrafish [32, 33]. In this study, 25% (6 out of 24) of tumors were

pPNETs in *sdp^{w12/+};rpL35^{hi258/+}*, while only 7% (2 out of 28) were pPNETs in *rpL35^{hi258/+}*. A pPNET is a type of round cell sarcoma belonging to the Ewing family of tumors, which also originate from the neural ectoderm where neural crest cells develop [42]. Considering a potential developmental role of *reck* in the zebrafish neural crest




Figure 2: *Reck* is a tumor suppressor gene in zebrafish MPNSTs. (A–B) Loss of the *reck* gene function facilitates tumorigenesis in either *rp* or *tp53* mutant background. Kaplan–Meier curves showing tumor-free survival of cohorts of single and double heterozygotes derived from *rpL35^{hi258/+}* x *sdp^{w12/+}* crosses (A) and *tp53^{M214K/+}* x *sdp^{w12/+}* crosses (B). Fish from all crosses were genotyped by PCR for each relevant mutation at 6–8 weeks of age, and housed segregated by genotype. In all panels, the numbers of fish (n) of each genotype are shown next to the genotype. The p values between the *rp* or *tp53* single heterozygote and the compound mutants are shown in the figures. The green line (*sdp^{w12/+}*) and black line (non-carrier) are overlapped in panel (A). (C) More zebrafish developed pPNETs in the *rpL35^{hi258/+};sdp^{w12/+}* compared to *rpL35^{hi258/+}*. Two-tailed fisher’s exact test was used for statistical analysis. $p = 0.008$. Asterisks indicate statistical significance (A–C). (D) A typical zebrafish with abdominal MNPST. The green arrow indicates the position of the tumor. (E) Hematoxylin and Eosin (H&E) staining showed typical histopathology of zebrafish MPNST, which is characterized by interlacing bundles and swirls of spindle shaped tumor cells. (F) A typical pPNET that was found in the tail region. The green arrow indicates the position of the tumor. (G) Representative examples of pPNET histology by H&E staining. The pseudorosettes (Flexner-Wintersteiner rosettes) are indicated with green arrows. (H) The wildtype allele is lost in some of the *tp53^{M214K/+};sdp^{w12/+}* tumors (tumor samples are labeled with asterisks). The *reck* gene was amplified by PCR using primers flanking the *sdp^{w12}* mutation site. The PCR product was then digested with *Ava* II, which only cuts the 233bps wildtype PCR product into 158bps and 75bps fragments [34]. T, tumor; N, normal tissue from caudal fin. Note, the lowest band is from the PCR primer polymerization. Scaler bar = 250 mm (D, F) Scale bar = 50 mm (E, G).

(reported previously [34] and Supplementary Figure 1), the loss of *reck* function is likely responsible for the tumor cell differentiation to pPNETs instead of MPNSTs. RECK was recently found to regulate WNT [43, 44] and NOTCH [45] signaling pathways, which are important for neural crest differentiation [46, 47] and MPNST tumorigenesis [48–50]. Future studies are required to reveal the detailed mechanism behind this phenomenon.

Identification of new TSGs is not only important for the understanding of cancer biology, but also the key for cancer targeted therapy. Cancer targeted therapy is generally believed to be aimed at an oncogene. However, TSGs are also important for cancer targeted therapy, since the discovery of synthetic lethality of the *PARP1* and *BRCA1/2* genes and their clinical applications using PARP inhibitors [51, 52]. Along the same direction,




Figure 3: Restoration of *RECK* in human MPNST cells suppresses cell growth. (A) *RECK* expression is lost or dramatically decreased in human MPNST cells (STS26T, 90-8TL, S462, T265T and MC2633) compared to a benign neurofibroma cell line, HEI193. (B) *RECK* expression was restored in lentiviral *RECK*-stable cells (STS26T, S462 and 90-8TL) induced by doxycycline (1 μ g/ml) shown by Western blots. (C, F, I) MTT assays revealed that cell growth was inhibited by *RECK* restoration in STS26T, S462 and 90-8TL cell lines, respectively. Growth curves were created using the optical density values collected over five days. (D, G, J) Independent cell growth by plate colony formation assays on STS26T, S462 and 90-8TL, respectively. Crystal violet stained cell colonies (diameter \geq 0.2 mm) were counted. Asterisk indicated the statistical significance by student *t*-test, $p < 0.05$. (E, H, K) Anchorage- independent growth by soft agar assay. The cells were seeded on agarose plates for colony formation assays, and colonies whose diameters are larger than 20 μ m were counted two weeks later. Asterisks indicated the statistical significance by student *t*-test, $p < 0.05$.

TSG restoration was also demonstrated as another way of targeted therapy based on the biology of TSGs [53]. In the case of the *RECK* gene, it is frequently reported to be silenced through promoter methylation in a variety of human cancers [54–59]. Thus, attempts were made to restore RECK expression using DNA methyltransferase (DNMT) inhibitor, 5-aza-2′-deoxycytidine (5-aza-dC), in colon cancer and salivary adenoid cystic carcinoma [60, 61]. Our cellular experiments on RECK restoration also support the possibility of RECK as a target for MPNST therapy. Furthermore, cancer driver genes that are located on the same chromosome region were found to cooperate and promote tumorigenesis [62]. The discovery of such combinations is very important in developing new cancer therapies. For example, this phenomenon has recently been reported in human colon cancers, where the *TP53* copy number loss was coupled with the loss of a neighboring gene, *POLR2A* [63]. Identification of other TSGs on the chromosome 9p might further help to understand the tumorigenic mechanism of RECK and to develop new cancer treatment strategies based on the concept of gene cooperation.

MATERIALS AND METHODS

Zebrafish lines, husbandry, and tumor onset analysis

Zebrafish were raised and maintained at the Purdue animal housing facility, which is approved by AAALAC. All experiments were carried out according to the protocols approved by the Purdue Animal Care and Use Committee (PACUC). The tumor-prone zebrafish lines carrying either the *tp53*^{M214K/M214K} point mutation, or insertional mutations in ribosomal protein gene, *rpL35*^{hi258/+}, have been described previously [32, 33]. The zebrafish *reck* point mutation carrier, *sdp*^{w12/+}, was acquired from the laboratory of Dr. David W. Raible at the University of Washington. After crossing, genotyping was carried out at 6–8 weeks of age using previously published genotyping methods [13, 34].

After genotyping, siblings of different genotypes were housed in adjacent tanks at similar densities to minimize environmental differences. From the sixth month, we started to check for fish with visible tumors. During tumorigenesis, fish were euthanized at first observation of tumors or other signs of illness. Any accidental fish loss was censored. The tumor tissue samples from the euthanized fish were processed for hematoxylin and eosin (H&E) staining. The tumor types were confirmed by a board certified veterinary pathologist, blinded to the experimental groups. The final Kaplan–Meier tumor free survival curves were generated, and log-rank tests were performed to determine statistical significance using GraphPad Prism 6.0 h [13].

RECK protein sequence and phylogeny analyses

RECK protein sequences were identified by a BLASTp search using the human RECK sequence as a query in Ensembl, NCBI, or JGI website. The longest sequence was preferentially chosen when there were multiple sequences. Multiple protein sequences (Supplementary Table 1) were aligned using MUSCLE program [64], and the FASTA format alignment can be found in the Supplementary File 1. The evolutionary model for phylogenetic analysis was identified using a best model test using maximum likelihood, and default parameters in MEGA6 [65]. A maximum likelihood phylogenetic tree was constructed using JTT + G with 1000 bootstrap replicates with PhyML 3.1 [66]. The final phylogenetic trees were viewed and generated with FigTree V1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree>). Gene intron-exon structures were analyzed using the longest transcripts in Ensembl.

Cell culture, stable cell lines, and cell growth assays

All experimental protocols using cell lines and plasmid constructs were approved by Purdue University institutional review board. The human MPNST cell lines were authenticated by ATCC using short tandem repeat profiling (April 25, 2016). Cells were cultured in DMEM with 10% heat inactivated fetal bovine serum, penicillin (100 IU/ml), and streptomycin (100 µg/ml). All cell cultures were carried out at 37° C in a humidified 5% CO₂ atmosphere. A full-length sequence of the *RECK* gene was amplified from HEK293T cells, and sequences were confirmed by Sanger sequencing. The *RECK* gene was subsequently cloned into pSlik-Neo vector (Addgene #25735). Stable cell establishment, MTT and plate colony assays were conducted following previously published methods [41]. For Western blot, commercial anti-RECK antibody (BD, #611512 or Cell Signaling #3433, 1:500 dilution) was used following an established method [41]. Statistical analyses were performed using GraphPad Prism 6.0 h. Data were analyzed using the un-paired student *t*-test. *p* < 0.05 was considered as statistically significant.

Soft agar assay

For each cell line, 1000 cells per well were seeded in six-well plates. 1.5 ml 1% agarose was plated at the bottom of plate. After solidification, 1 ml of single-cell suspension in 0.7% agarose in DMEM supplemented with 10% FBS were added in each well. Then, the plates were incubated at 37° C in a humidified incubator with 5% CO₂. On the second day, doxycycline (1 µg/ml) or PBS (phosphate-buffered saline) were added to induce RECK expression. The medium was refreshed every 2 days to maintain

constant expression of RECK. Two weeks later, cells were stained with 0.01% crystal violet solution. After multiple washes with PBS, the cell colonies were visualized and imaged under a bright field microscope. Six sub-images at different focus in the same field were captured at 40× magnification, and all the images of the same field were Z stacked using Image J software. Ten random bright field images of each sample were captured. Total number of formed foci (diameter > 20 μm) in each field were counted manually in Image J software. Three technical replicates were included, and each experiment was repeated at least three times for each experimental condition.

***In situ* hybridization**

The primers Dr.reck-97F (5' CACCATGAGCGG GTGTCTCCAGATCCTCA3') and Dr.reck-2961R (5' GAGGTCAGAGGTCAGGGTCAGGAT3') were used for the zebrafish *reck* gene cloning PCR. The zebrafish *reck* gene cloning and *In Situ* hybridization were performed using the standard methods as previously described [67].

Author contributions

GZ designed the project and coordinated research; RK, MRS, ANV, and GZ performed experiments; RK, MRS, YLJ, JYY, SKM, and GZ analyzed the results; GZ wrote the original draft. MRS, RK, YLJ, JYY, SKM, and GZ reviewed and edited the manuscript. All authors read and approved the final manuscript.

ACKNOWLEDGMENTS AND FUNDING

We gratefully acknowledge the Heyward Foundation for supporting our research. We thank Dr. David W. Raible for sharing the *sdp^{w12}* fish. We also thank Dr. George De Vries, and Dr. Karen Cichowski for kindly providing us the MPNST cell lines. We thank Guolin Chai and Xinmin Yan for technical help generating the pSlik-RECK construct, and Zhibin Cui and Pei-Chieh Tien for cell line and antibody validations. The authors also thank the support from the Purdue University Center for Cancer Research, NIH grant P30 CA023168, and Jim and Diann Robbers Cancer Research Grant for New Investigators. We thank the Purdue Histology Research Laboratory for preparation of tissue sections.

CONFLICTS OF INTEREST

None.

REFERENCES

1. Ducatman BS, Scheithauer BW, Piepgras DG, Reiman HM, Ilstrup DM. Malignant peripheral nerve sheath tumors.

A clinicopathologic study of 120 cases. *Cancer*. 1986; 57:2006–21.

2. Kolberg M, Holand M, Agesen TH, Brekke HR, Liestol K, Hall KS, Mertens F, Picci P, Smeland S, Lothe RA. Survival meta-analyses for >1800 malignant peripheral nerve sheath tumor patients with and without neurofibromatosis type 1. *Neuro Oncol*. 2013; 15:135–47. <https://doi.org/10.1093/neuonc/nos287>.
3. Durbin AD, Ki DH, He S, Look AT. Malignant Peripheral Nerve Sheath Tumors. *Adv Exp Med Biol*. 2016; 916:495–530. https://doi.org/10.1007/978-3-319-30654-4_22.
4. Piotrowski A, Xie J, Liu YF, Poplawski AB, Gomes AR, Madanecki P, Fu C, Crowley MR, Crossman DK, Armstrong L, Babovic-Vuksanovic D, Bergner A, Blakeley JO, et al. Germline loss-of-function mutations in LZTR1 predispose to an inherited disorder of multiple schwannomas. *Nat Genet*. 2014; 46:182–87. <https://doi.org/10.1038/ng.2855>.
5. McClatchey AI. Neurofibromatosis. *Annu Rev Pathol*. 2007; 2:191–216. <https://doi.org/10.1146/annurev.pathol.2.010506.091940>.
6. Farid M, Demicco EG, Garcia R, Ahn L, Merola PR, Cioffi A, Maki RG. Malignant peripheral nerve sheath tumors. *Oncologist*. 2014; 19:193–201. <https://doi.org/10.1634/theoncologist.2013-0328>.
7. Stratton MR, Campbell PJ, Futreal PA. The cancer genome. *Nature*. 2009; 458:719–24. <https://doi.org/10.1038/nature07943>.
8. Weaver BA, Cleveland DW. Does aneuploidy cause cancer? *Curr Opin Cell Biol*. 2006; 18:658–67.
9. Kim M, Gans JD, Nogueira C, Wang A, Paik JH, Feng B, Brennan C, Hahn WC, Cordon-Cardo C, Wagner SN, Flotte TJ, Duncan LM, Granter SR, et al. Comparative oncogenomics identifies NEDD9 as a melanoma metastasis gene. *Cell*. 2006; 125:1269–81. <https://doi.org/10.1016/j.cell.2006.06.008>.
10. Maser RS, Choudhury B, Campbell PJ, Feng B, Wong KK, Protopopov A, O'Neil J, Gutierrez A, Ivanova E, Perna I, Lin E, Mani V, Jiang S, et al. Chromosomally unstable mouse tumours have genomic alterations similar to diverse human cancers. *Nature*. 2007; 447:966–71.
11. Zender L, Spector MS, Xue W, Flemming P, Cordon-Cardo C, Silke J, Fan ST, Luk JM, Wigler M, Hannon GJ, Mu D, Lucito R, Powers S, et al. Identification and validation of oncogenes in liver cancer using an integrative oncogenomic approach. *Cell*. 2006; 125:1253–67. <https://doi.org/10.1016/j.cell.2006.05.030>.
12. Tang J, Le S, Sun L, Yan X, Zhang M, Macleod J, Leroy B, Northrup N, Ellis A, Yeatman TJ, Liang Y, Zwick ME, Zhao S. Copy number abnormalities in sporadic canine colorectal cancers. *Genome Res*. 2010; 20:341–50. <https://doi.org/10.1101/gr.092726.109>.
13. Zhang G, Hoersch S, Amsterdam A, Whittaker CA, Beert E, Catchen JM, Farrington S, Postlethwait JH, Legius E, Hopkins N, Lees JA. Comparative oncogenomic analysis of copy number alterations in human and zebrafish tumors

- enables cancer driver discovery. *PLoS Genet.* 2013; 9:e1003734. <https://doi.org/10.1371/journal.pgen.1003734>.
14. Zhang G, Hoersch S, Amsterdam A, Whittaker CA, Lees JA, Hopkins N. Highly aneuploid zebrafish malignant peripheral nerve sheath tumors have genetic alterations similar to human cancers. *Proc Natl Acad Sci U S A.* 2010; 107:16940–5. <https://doi.org/10.1073/pnas.1011548107>.
 15. Chen EY, Dobrinski KP, Brown KH, Clagg R, Edelman E, Ignatius MS, Chen JY, Brockmann J, Nielsen GP, Ramaswamy S, Keller C, Lee C, Langenau DM. Cross-species array comparative genomic hybridization identifies novel oncogenic events in zebrafish and human embryonal rhabdomyosarcoma. *PLoS Genet.* 2013; 9:e1003727. <https://doi.org/10.1371/journal.pgen.1003727>.
 16. White R, Rose K, Zon L. Zebrafish cancer: the state of the art and the path forward. *Nat Rev Cancer.* 2013; 13:624–36. <https://doi.org/10.1038/nrc3589>.
 17. Liu S, Leach SD. Zebrafish models for cancer. *Annu Rev Pathol.* 2011; 6:71–93. <https://doi.org/10.1146/annurev-pathol-011110-130330>.
 18. Gunawan B, Schulten HJ, von Heydebreck A, Schmidt B, Enders C, Hoer J, Langer C, Schuler P, Schindler CG, Kuhlitz J, Fuzesi L. Site-independent prognostic value of chromosome 9q loss in primary gastrointestinal stromal tumours. *J Pathol.* 2004; 202:421–9. <https://doi.org/10.1002/path.1537>.
 19. Lichun Y, Ching Tang CM, Wai Lau K, Lung ML. Frequent loss of heterozygosity on chromosome 9 in Chinese esophageal squamous cell carcinomas. *Cancer Lett.* 2004; 203:71–7.
 20. Van Tilborg AA, Hekman AC, Vissers KJ, van der Kwast TH, Zwarthoff EC. Loss of heterozygosity on chromosome 9 and loss of chromosome 9 copy number are separate events in the pathogenesis of transitional cell carcinoma of the bladder. *Int J Cancer.* 1998; 75:9–14.
 21. Zhu Y, Spitz MR, Strom S, Tomlinson GE, Amos CI, Minna JD, Wu X. A case-control analysis of lymphocytic chromosome 9 aberrations in lung cancer. *Int J Cancer.* 2002; 102:536–40. <https://doi.org/10.1002/ijc.10762>.
 22. Huntly BJ, Bench AJ, Delabesse E, Reid AG, Li J, Scott MA, Campbell L, Byrne J, Pinto E, Brizard A, Niedermeiser D, Nacheva EP, Guilhot F, et al. Derivative chromosome 9 deletions in chronic myeloid leukemia: poor prognosis is not associated with loss of ABL-BCR expression, elevated BCR-ABL levels, or karyotypic instability. *Blood.* 2002; 99:4547–53.
 23. Eiriksdottir G, Sigurdsson A, Jonasson JG, Agnarsson BA, Sigurdsson H, Gudmundsson J, Bergthorsson JT, Barkardottir RB, Egilsson V, Ingvarsson S. Loss of heterozygosity on chromosome 9 in human breast cancer: association with clinical variables and genetic changes at other chromosome regions. *Int J Cancer.* 1995; 64:378–82.
 24. Beroukhi R, Mermel CH, Porter D, Wei G, Raychaudhuri S, Donovan J, Barretina J, Boehm JS, Dobson J, Urashima M, Mc Henry KT, Pinchback RM, Ligon AH, et al. The landscape of somatic copy-number alteration across human cancers. *Nature.* 2010; 463:899–905. <https://doi.org/10.1038/nature08822>.
 25. Beert E, Brems H, Daniels B, De Wever I, Van Calenbergh F, Schoenaers J, Debiec-Rychter M, Gevaert O, De Raedt T, Van Den Bruel A, de Ravel T, Cichowski K, Kluwe L, et al. Atypical neurofibromas in neurofibromatosis type 1 are premalignant tumors. *Genes Chromosomes Cancer.* 2011; 50:1021–32. <https://doi.org/10.1002/gcc.20921>.
 26. Noda M, Takahashi C. Recklessness as a hallmark of aggressive cancer. *Cancer Sci.* 2007; 98:1659–65. <https://doi.org/10.1111/j.1349-7006.2007.00588.x>.
 27. Takahashi C, Sheng Z, Horan TP, Kitayama H, Maki M, Hitomi K, Kitaura Y, Takai S, Sasahara RM, Horimoto A, Ikawa Y, Ratzkin BJ, Arakawa T, et al. Regulation of matrix metalloproteinase-9 and inhibition of tumor invasion by the membrane-anchored glycoprotein RECK. *Proc Natl Acad Sci U S A.* 1998; 95:13221–6.
 28. Alexius-Lindgren M, Andersson E, Lindstedt I, Engstrom W. The RECK gene and biological malignancy—its significance in angiogenesis and inhibition of matrix metalloproteinases. *Anticancer Res.* 2014; 34:3867–73.
 29. Jacomaso T, Trombetta-Lima M, Sogayar MC, Winnischofer SM. Downregulation of reversion-inducing cysteine-rich protein with Kazal motifs in malignant melanoma: inverse correlation with membrane-type 1-matrix metalloproteinase and tissue inhibitor of metalloproteinase 2. *Melanoma Res.* 2014; 24:32–9. <https://doi.org/10.1097/CMR.0000000000000039>.
 30. Span PN, Sweep CG, Manders P, Beex LV, Leppert D, Lindberg RL. Matrix metalloproteinase inhibitor reversion-inducing cysteine-rich protein with Kazal motifs: a prognostic marker for good clinical outcome in human breast carcinoma. *Cancer.* 2003; 97:2710–5. <https://doi.org/10.1002/cncr.11395>.
 31. Oh J, Takahashi R, Kondo S, Mizoguchi A, Adachi E, Sasahara RM, Nishimura S, Imamura Y, Kitayama H, Alexander DB, Ide C, Horan TP, Arakawa T, et al. The membrane-anchored MMP inhibitor RECK is a key regulator of extracellular matrix integrity and angiogenesis. *Cell.* 2001; 107:789–800.
 32. Berghmans S, Murphey RD, Wienholds E, Neubergh D, Kutok JL, Fletcher CD, Morris JP, Liu TX, Schulte-Merker S, Kanki JP, Plasterk R, Zon LI, Look AT. tp53 mutant zebrafish develop malignant peripheral nerve sheath tumors. *Proc Natl Acad Sci U S A.* 2005; 102:407–12. <https://doi.org/10.1073/pnas.0406252102>.
 33. Amsterdam A, Sadler KC, Lai K, Farrington S, Bronson RT, Lees JA, Hopkins N. Many ribosomal protein genes are cancer genes in zebrafish. *PLoS Biol.* 2004; 2:E139. <https://doi.org/10.1371/journal.pbio.0020139>.
 34. Prendergast A, Linbo TH, Swarts T, Ungos JM, McGraw HF, Krispin S, Weinstein BM, Raible DW. The

- metalloproteinase inhibitor Reck is essential for zebrafish DRG development. *Development*. 2012; 139:1141–52. <https://doi.org/10.1242/dev.072439>.
35. Lai K, Amsterdam A, Farrington S, Bronson RT, Hopkins N, Lees JA. Many ribosomal protein mutations are associated with growth impairment and tumor predisposition in zebrafish. *Dev Dyn*. 2009; 238:76–85. <https://doi.org/10.1002/dvdy.21815>.
 36. MacInnes AW, Amsterdam A, Whittaker CA, Hopkins N, Lees JA. Loss of p53 synthesis in zebrafish tumors with ribosomal protein gene mutations. *Proc Natl Acad Sci USA*. 2008; 105:10408–13. <https://doi.org/10.1073/pnas.0805036105>.
 37. Berger AH, Knudson AG, Pandolfi PP. A continuum model for tumour suppression. *Nature*. 2011; 476:163–9. <https://doi.org/10.1038/nature10275>.
 38. Ventura A, Kirsch DG, McLaughlin ME, Tuveson DA, Grimm J, Lintault L, Newman J, Reczek EE, Weissleder R, Jacks T. Restoration of p53 function leads to tumour regression *in vivo*. *Nature*. 2007; 445:661–5. <https://doi.org/10.1038/nature05541>.
 39. Miller SJ, Rangwala F, Williams J, Ackerman P, Kong S, Jegga AG, Kaiser S, Aronow BJ, Frahm S, Kluge L, Mautner V, Upadhyaya M, Muir D, et al. Large-scale molecular comparison of human schwann cells to malignant peripheral nerve sheath tumor cell lines and tissues. *Cancer Res*. 2006; 66:2584–91. <https://doi.org/10.1158/0008-5472.can-05-3330>.
 40. De Raedt T, Beert E, Pasmant E, Luscan A, Brems H, Ortonne N, Helin K, Hornick JL, Mautner V, Kehrer-Sawatzki H, Clapp W, Bradner J, Vidaud M, et al. PRC2 loss amplifies Ras-driven transcription and confers sensitivity to BRD4-based therapies. *Nature*. 2014; 514:247–51. <https://doi.org/10.1038/nature13561>.
 41. Cui Z, Shen Y, Chen KH, Mittal SK, Yang JY, Zhang G. KANK1 inhibits cell growth by inducing apoptosis through regulating CXXC5 in human malignant peripheral nerve sheath tumors. *Sci Rep*. 2017; 7:40325. <https://doi.org/10.1038/srep40325>.
 42. Tsokos M, Alaggio RD, Dehner LP, Dickman PS. Ewing sarcoma/peripheral primitive neuroectodermal tumor and related tumors. *Pediatr Dev Pathol*. 2012; 15:108–26. <https://doi.org/10.2350/11-08-1078-PB.1>.
 43. Ulrich F, Carretero-Ortega J, Menendez J, Narvaez C, Sun B, Lancaster E, Pershad V, Trzaska S, Veliz E, Kamei M, Prendergast A, Kidd KR, Shaw KM, et al. Reck enables cerebrovascular development by promoting canonical Wnt signaling. *Development*. 2016; 143:1055. <https://doi.org/10.1242/dev.136507>.
 44. Cho C, Smallwood PM, Nathans J. Reck and Gpr124 Are Essential Receptor Cofactors for Wnt7a/Wnt7b-Specific Signaling in Mammalian CNS Angiogenesis and Blood-Brain Barrier Regulation. *Neuron*. 2017; 95:1221–5. <https://doi.org/10.1016/j.neuron.2017.08.032>.
 45. Park S, Lee C, Sabharwal P, Zhang M, Meyers CL, Sockanathan S. GDE2 promotes neurogenesis by glycosylphosphatidylinositol-anchor cleavage of RECK. *Science*. 2013; 339:324–8. <https://doi.org/10.1126/science.1231921>.
 46. Dorsky RI, Moon RT, Raible DW. Control of neural crest cell fate by the Wnt signalling pathway. *Nature*. 1998; 396:370–3. <https://doi.org/10.1038/24620>.
 47. Cornell R, Eisen J. Notch in the pathway: The roles of Notch signaling in neural crest development. *Semin Cell Dev Biol*. 2005; 16:663–72. <https://doi.org/10.1016/j.semcdb.2005.06.009>.
 48. Muraguchi T, Takegami Y, Ohtsuka T, Kitajima S, Chandana EP, Omura A, Miki T, Takahashi R, Matsumoto N, Ludwig A, Noda M, Takahashi C. RECK modulates Notch signaling during cortical neurogenesis by regulating ADAM10 activity. *Nat Neurosci*. 2007; 10:838–45. <https://doi.org/10.1038/nn1922>.
 49. Noda M, Vallon M, Kuo CJ. The Wnt7's Tale: A story of an orphan who finds her tie to a famous family. *Cancer Sci*. 2016; 107:576–82. <https://doi.org/10.1111/cas.12924>.
 50. Watson AL, Rahrmann EP, Moriarity BS, Choi K, Conboy CB, Greeley AD, Halfond AL, Anderson LK, Wahl BR, Keng VW, Rizzardi AE, Forster CL, Collins MH, et al. Canonical Wnt/beta-catenin Signaling Drives Human Schwann Cell Transformation, Progression, and Tumor Maintenance. *Cancer Discov*. 2013; 3:674–89. <https://doi.org/10.1158/2159-8290.CD-13-0081>.
 51. Morris LG, Chan TA. Therapeutic targeting of tumor suppressor genes. *Cancer*. 2015; 121:1357–68. <https://doi.org/10.1002/cncr.29140>.
 52. Wang H, Han H, Mousses S, Von Hoff DD. Targeting loss-of-function mutations in tumor-suppressor genes as a strategy for development of cancer therapeutic agents. *Semin Oncol*. 2006; 33:513–20. <https://doi.org/10.1053/j.seminoncol.2006.04.013>.
 53. Xue W, Zender L, Miething C, Dickins RA, Hernando E, Krizhanovsky V, Cordon-Cardo C, Lowe SW. Senescence and tumour clearance is triggered by p53 restoration in murine liver carcinomas. *Nature*. 2007; 445:656–60. <https://doi.org/10.1038/nature05529>.
 54. Du YY, Dai DQ, Yang Z. Role of RECK methylation in gastric cancer and its clinical significance. *World J Gastroenterol*. 2010; 16:904–8.
 55. Pesta M, Kulda V, Topolcan O, Safranek J, Vrzalova J, Cerny R, Holubec L. Significance of methylation status and the expression of RECK mRNA in lung tissue of patients with NSCLC. *Anticancer Res*. 2009; 29:4535–9.
 56. Lu XX, Zhang SM, Fang Y, Wang ZT, Xie JJ, Zhan Q, Deng XX, Chen H, Jin JB, Peng CH, Li HW, Shen BY. Clinical significance of RECK promoter methylation in pancreatic ductal adenocarcinoma. *Tumour Biol*. 2013; 34:3339–43. <https://doi.org/10.1007/s13277-013-0903-z>.
 57. Zhang CS, Ling Y, Zhang CH, Xu Y, Gao L, Li R, Zhu J, Fan LY, Wei LX. The Silencing of RECK Gene is Associated with Promoter Hypermethylation and Poor

- Survival in Hepatocellular Carcinoma. *Int J Biol Sci.* 2012; 8:451–8. <https://doi.org/10.7150/ijbs.4038>.
58. Chang HC, Cho CY, Hung WC. Downregulation of RECK by promoter methylation correlates with lymph node metastasis in non-small cell lung cancer. *Cancer Sci.* 2007; 98:169–73. <https://doi.org/10.1111/j.1349-7006.2006.00367.x>.
 59. Shi G, Yoshida Y, Yuki K, Nishimura T, Kawata Y, Kawashima M, Iwaisako K, Yoshikawa K, Kurebayashi J, Toi M, Noda M. Pattern of RECK CpG methylation as a potential marker for predicting breast cancer prognosis and drug-sensitivity. *Oncotarget.* 2016; 7:82158–69. <https://doi.org/10.18632/oncotarget.8620>.
 60. Zhou XQ, Huang SY, Zhang DS, Zhang SZ, Li WG, Chen ZW, Wu HW. Effects of 5-aza-2'-deoxycytidine on RECK gene expression and tumor invasion in salivary adenoid cystic carcinoma. *Braz J Med Biol Res.* 2015; 48:254–60. <https://doi.org/10.1590/1414-431X20144102>.
 61. Cho CY, Wang JH, Chang HC, Chang CK, Hung WC. Epigenetic inactivation of the metastasis suppressor RECK enhances invasion of human colon cancer cells. *J Cell Physiol.* 2007; 213:65–9. <https://doi.org/10.1002/jcp.21089>.
 62. Xue W, Kitzing T, Roessler S, Zuber J, Krasnitz A, Schultz N, Revill K, Weissmueller S, Rappaport AR, Simon J, Zhang J, Luo W, Hicks J, et al. A cluster of cooperating tumor-suppressor gene candidates in chromosomal deletions. *Proc Natl Acad Sci U S A.* 2012; 109:8212–7. <https://doi.org/10.1073/pnas.1206062109>.
 63. Liu Y, Zhang X, Han C, Wan G, Huang X, Ivan C, Jiang D, Rodriguez-Aguayo C, Lopez-Berestein G, Rao PH, Maru DM, Pahl A, He X, et al. TP53 loss creates therapeutic vulnerability in colorectal cancer. *Nature.* 2015; 520:697–701. <https://doi.org/10.1038/nature14418>.
 64. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 2004; 32:1792–7. <https://doi.org/10.1093/nar/gkh340>.
 65. Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol.* 2013; 30:2725–9. <https://doi.org/10.1093/molbev/mst197>.
 66. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol.* 2010; 59:307–21. <https://doi.org/10.1093/sysbio/syq010>.
 67. Hensley MR, Cui Z, Chua RFM, Simpson S, Shammass NL, Yang JY, Leung YF, Zhang G. Evolutionary and developmental analysis reveals KANK genes were co-opted for vertebrate vascular development. *Sci Rep.* 2016; 6:27816. <https://doi.org/10.1038/srep27816>.