

Correction: Long non-coding RNAs could act as vectors for paternal heredity of high fat diet-induced obesity

Tian An^{1,*}, Teng Zhang^{2,*}, Fei Teng¹, Jia-Cheng Zuo¹, Yan-Yun Pan¹, Yu-Fei Liu³, Jia-Nan Miao¹, Yu-Jie Gu¹, Na Yu¹, Dan-Dan Zhao¹, Fang-Fang Mo¹, Si-Hua Gao¹ and Guangjian Jiang¹

¹ Diabetes Research Center, Beijing University of Chinese Medicine, Beijing, China

² State Key Laboratory of Stem Cell and Reproductive Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing, China

³ The Third Affiliated Hospital, Beijing University of Chinese Medicine, Beijing, China

* These authors have contributed equally to this work

Published: January 22, 2019

Copyright: An et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been corrected: The correct GEO number information is given below:

Data acquisition

Slides were scanned with the Microarray Scanner (cat. no. G2565CA; Agilent Technologies) using the following default settings: dye channel = green, scan resolution = 3 μ m, PMT = 100%, and 20 bit. Data were extracted with Feature Extraction v.10.7 (Agilent Technologies). Raw data were normalised with the quantile algorithm of GeneSpring v.12.6.1 (Agilent Technologies). The microarray data presented in this article have been deposited in the National Center for Biotechnology Information Gene Expression Omnibus (GEO) and are accessible through GEO Series accession number GSE118039.

Original article: Oncotarget. 2017; 8:47876-47889. <https://doi.org/10.18632/oncotarget.18138>