Erratum

Erratum: Predicting clinically significant prostate cancer using DCE-MRI habitat descriptors

N. Andres Parra[1], Hong Lu[1,2], Qian Li[2], Radka Stoyanova[3], Alan Pollack[3], Sanoj Punnen[4], Jung Choi[5], Mahmoud Abdalah[1], Christopher Lopez[3], Kenneth Gage[5], Jong Y. Park[6], Yamoah Kosj[6,7], Julio M. Pow-Sang[8], Robert J. Gillies[1,5] and Yoganand Balagurunathan[1]

[1] Department of Cancer Physiology, H.L. Moffitt Cancer Center, Tampa, FL, USA

[2] Department of Radiology, Tianjin Medical University Cancer Institute and Hospital, Tianjin, China

[3] Department of Radiation Oncology, University of Miami Miller School of Medicine, Miami, FL, USA

[4] Department of Urology, University of Miami Miller School of Medicine, Miami, FL, USA

[5] Department of Radiology, H.L. Moffitt Cancer Center, Tampa, FL, USA

[6] Department of Cancer Epidemiology, H.L. Moffitt Cancer Center, Tampa, FL, USA

[7] Department of Radiation Oncology, H.L. Moffitt Cancer Center, Tampa, FL, USA

[8] Department of Urology, H.L. Moffitt Cancer Center, Tampa, FL, USA

Published: March 12, 2019

This article has been corrected: During production, the author name for this article was given incorrectly. The proper author name is shown below.

N. Andres Parra

Original article: Oncotarget. 2018; 9:37125-37136. https://doi.org/10.18632/oncotarget.26437