

Correction: Nucleolin represses transcription of the androgen receptor gene through a G-quadruplex

Cindy K. Miranti^{1,2}, Sara Moore³, Yongeun Kim⁴, Venkateshwar Reddy Chappeta⁴, Kui Wu⁴, Biswanath De⁴, Vijay Gokhale⁵, Laurence H. Hurley^{1,4,5} and Elsa M. Reyes-Reyes^{1,2,3}

¹University of Arizona Cancer Center, University of Arizona, Tucson, AZ 85724, USA

²Department of Cellular and Molecular Medicine, Tucson, AZ 85724, USA

³University of Arizona College of Medicine, Division of Pulmonary, Allergy, Critical Care, and Sleep Medicine, Tucson, AZ 85724, USA

⁴College of Pharmacy, University of Arizona, Tucson, AZ 85724, USA

⁵BIO5 Institute, University of Arizona, Tucson, AZ 85724, USA

Published: June 30, 2020

Copyright: Miranti et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been corrected: The Acknowledgements section has been updated as follows:

ACKNOWLEDGMENTS

These studies were supported by funds from IRG-16-124-37 from the American Cancer Society and University of Arizona Career Development Award to EMRR and by NIH award R01CA177585 to L.H.H.

Original article: Oncotarget. 2020; 11:1758–1776. <https://doi.org/10.18632/oncotarget.27589>