

Correction: MiR-101 reverses the hypomethylation of the LMO3 promoter in glioma cells

Xiaoping Liu^{1,2,*}, Qianqian Lei^{1,*}, Zhibin Yu¹, Gang Xu^{1,4}, Hailin Tang², Wei Wang¹, Zeyou Wang¹, Guiyuan Li^{3,1} and Minghua Wu^{3,1}

¹Hunan Provincial Tumor Hospital and the Affiliated Tumor Hospital of Xiangya Medical School, Central South University, Changsha 410013, Hunan, China

²Department of Breast Oncology, Sun Yat-Sen University Cancer Center, State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, Guangzhou 510060, Guangdong, China

³School of Basic Medical Science, Cancer Research Institute, Central South University, Key Laboratory of Carcinogenesis and Cancer Invasion, Ministry of Education, Key Laboratory of Carcinogenesis, Ministry of Health, Changsha 410078, Hunan, China

⁴Medical College, University of South China, Hengyang 421001, Hunan, China

*These authors have contributed equally to this work

Published: December 08, 2020

Copyright: Liu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been corrected: During assembly of Figure 5A, the authors made an error in the vector and scramble sequences co-transfected group, whose picture was the same as the si-NC and scramble sequences co-transfected group. The corrected Figure 5A is shown below. The authors declare that these corrections do not change the results or conclusions of this paper.

Original article: Oncotarget. 2015; 6:7930–7943. <https://doi.org/10.18632/oncotarget.3181>




Figure 5: MiR-101 induces U251 cell apoptosis via LMO3. (A) LMO3 overexpression counteracted miR-101-promoted apoptosis ($\times 100$). Left: Hoechst 33258 staining was used to detect the level of apoptosis in U251 cells.