

Retraction

Retraction: Hypoxia-inducible factor 1a induces phenotype switch of human aortic vascular smooth muscle cell through PI3K/AKT/AEG-1 signaling

Kai Liu¹, Changcun Fang¹, Yuwen Shen¹, Zhengqin Liu¹, Min Zhang¹, Bingbing Ma¹ and Xinyan Pang¹

¹Department of Cardiovascular Surgery, Qilu Hospital of Shandong University, Jinan 250012, Shandong, China

Published: June 20, 2024

Copyright: © 2024 Liu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#) (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been retracted: Multiple instances of image duplication were discovered in this paper. Specifically, the GAPDH blots in 1st and 2nd panels of Figure 1A are internal duplicates even though the expression of HIF-1 α mRNA was measured at different conditions. Additionally, multiple blots in Figure 3A and Figure 8, panels A and B, are duplicates of those in Figure 4B and 4C of an earlier published paper unrelated by subject [1]. Therefore, with the agreement of all authors, the Scientific Integrity office at Oncotarget has decided to retract this paper.

Original article: Oncotarget. 2017; 8:33343–33352. <https://doi.org/10.18632/oncotarget.16448>

REFERENCES

1. Yuan Z, Chen D, Chen X, Yang H, Wei Y. Overexpression of trefoil factor 3 (TFF3) contributes to the malignant progression in cervical cancer cells. *Cancer Cell Int.* 2017; 17:7. <https://doi.org/10.1186/s12935-016-0379-1>. [PubMed]